

## Rapid Assessment Reference Condition Model

The Rapid Assessment is a component of the LANDFIRE project. Reference condition models for the Rapid Assessment were created through a series of expert workshops and a peer-review process in 2004-2005. For more information, please visit [www.landfire.gov](http://www.landfire.gov). Please direct questions to [helpdesk@landfire.gov](mailto:helpdesk@landfire.gov).

### Potential Natural Vegetation Group (PNVG):

R5SOFPrf

Southern Floodplain - Rare Fire

### General Information

**Contributors** (additional contributors may be listed under "Model Evolution and Comments")

#### Modelers

Bruce Davenport      bdavenport@fs.fed.us  
Kevin Robertson      kevin@ttrs.org

#### Reviewers

Doug Zollner      dzollner@tnc.org  
Maria Melnechuk      maria\_melnechuk@tnc.org

#### Vegetation Type

Forested

#### Dominant Species\*

TADI2      ACRU  
TAAS      FRPE  
NYAQ2      SANI  
QULY

#### General Model Sources

- Literature  
 Local Data  
 Expert Estimate

#### LANDFIRE Mapping Zones

45      58  
46      55  
37

#### Rapid Assessment Model Zones

- California       Pacific Northwest  
 Great Basin       South Central  
 Great Lakes       Southeast  
 Northeast       S. Appalachians  
 Northern Plains       Southwest  
 N-Cent.Rockies

#### **Geographic Range**

This PNVG occurs from east Texas to Virginia within the Coastal Plain and lower Piedmont and up the Mississippi River basin to southern Illinois.

#### **Biophysical Site Description**

The landscape includes sloughs and abandoned channels which are flooded most or all of a given year as well as backswamps and depressions within the flood plain which are frequently flooded and where soils remain saturated or with water table close to the surface much of the year.

#### **Vegetation Description**

The vegetation is generally close canopied forests ranging from standing water to floodplain depressions. The canopy is normally dominated by cypress and tupelo under the wettest conditions and overcup oak or maple and ash on the drier end.

#### **Disturbance Description**

Weather, primarily wind and flooding, is the dominant disturbance agent in this type and includes wind damage from hurricanes and tornadoes as well as scouring, changing streamcourses, and inundated young stands. Because of its moisture regime, fire is rare, occurring only during extreme drought conditions. In addition, replacement fire requires not only extended drought but accumulated fuel by drift or deep "duff" development (may be normally submerged). Insect outbreaks would occur infrequently in closed canopy states.

#### **Adjacency or Identification Concerns**

Although the Southern Floodplain Forest is included in both Kuchler and coarse scale, a difference in hydroperiod from other, adjacent types often results in a dependence on drought for regeneration and

\*Dominant and Indicator Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

subsequently, stand structure. In the absence of characteristic vegetation, the break in moisture gradient between the drier end of this type and even drier types may be unclear.

**Scale Description**

**Sources of Scale Data**  Literature  Local Data  Expert Estimate

The landscape has adequate coverage to encompass natural variation. At either end of the spectrum, large swamps may cover millions of acres (Atchafalaya) while individual oxbows may be less than one hundred.

**Issues/Problems**

Contains long-lived species with very long fire return interval and, often, uncommon conditions required to complete life history. Literature and documentation of modeled conditions, especially fire, are not readily available.

**Model Evolution and Comments**

Suggested reviewers: Tom Foti (tom@arkansasheritage.org), Paul Hamel (phamel@fs.fed.us), Charles Klimas (Waterways Exp. Sta.), Rob Evans (? Formerly NatureServe). Insect/disease was added as a disturbance after peer review.

**Succession Classes**  
*Succession classes are the equivalent of "Vegetation Fuel Classes" as defined in the Interagency FRCC Guidebook (www.frcc.gov).*

**Class A 10%**

Early1 All Structures

**Description**

0-19 years. Seedlings, saplings, and some sprouts on drier sites, in openings created by flood scouring, changed streamcourses, wind damage, or, infrequently, fire. Primarily composed of major overstory species with transient herbaceous plants and shrub, small trees and woody vines; the latter, woody group occurring more often on drier sites.

**Indicator Species\* and Canopy Position**

TADI2 Upper  
 NYAQ2 Upper  
 QULY Upper  
 FRPE Upper

**Upper Layer Lifeform**

- Herbaceous
- Shrub
- Tree

**Fuel Model 9**

**Structure Data (for upper layer lifeform)**

	<i>Min</i>	<i>Max</i>
<i>Cover</i>	40 %	80 %
<i>Height</i>	Tree Regen <5m	Tree Short 5-9m
<i>Tree Size Class</i>	Sapling >4.5ft; <5"DBH	

Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

**Class B 25%**

Mid1 Closed

**Description**

20-99 years. Dominated by young to early mature canopy species with a few obligate midstory species on less frequently flooded sites. Longer hydroperiod sites at least seasonally flooded and typically display a single, closed canopy layer.

**Indicator Species\* and Canopy Position**

TADI2 Upper  
 NYAQ2 Upper  
 QULY Upper  
 FRPE Upper

**Upper Layer Lifeform**

- Herbaceous
- Shrub
- Tree

**Fuel Model 9**

**Structure Data (for upper layer lifeform)**

	<i>Min</i>	<i>Max</i>
<i>Cover</i>	70 %	100 %
<i>Height</i>	Tree Short 5-9m	Tree Tall 25-49m
<i>Tree Size Class</i>	Medium 9-21"DBH	

Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

\*Dominant and Indicator Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

**Class C 20%**

Late1 Open  
**Description**  
100+ years. Early to, more often, late mature open canopy in long-term flooded conditions. Created during wet periods that prevent replacement of mortality.

**Indicator Species\* and Canopy Position**

TADI2 Upper  
NYAQ2 Upper  
QULY Upper  
FRPE Upper

**Upper Layer Lifeform**

- Herbaceous
- Shrub
- Tree

**Fuel Model** 9

**Structure Data (for upper layer lifeform)**

	Min	Max
Cover	15 %	60 %
Height	Tree Medium 10-24m	Tree Tall 25-49m
Tree Size Class	Large 21-33"DBH	

Upper layer lifeform differs from dominant lifeform.  
Height and cover of dominant lifeform are:

**Class D 45%**

Late1 Closed  
**Description**  
100+ years. Early to late mature closed canopy generally occurring as a single overstory layer, particularly on wetter sites. Drier sites will contain some midstory and young overstory species.

**Indicator Species\* and Canopy Position**

TADI2 Upper  
NYAQ2 Upper  
QULY Upper  
FRPE Upper

**Upper Layer Lifeform**

- Herbaceous
- Shrub
- Tree

**Fuel Model** 9

**Structure Data (for upper layer lifeform)**

	Min	Max
Cover	60 %	90 %
Height	Tree Medium 10-24m	Tree Tall 25-49m
Tree Size Class	Large 21-33"DBH	

Upper layer lifeform differs from dominant lifeform.  
Height and cover of dominant lifeform are:

**Class E 0%**

Late1 All Structures  
**Description**

**Indicator Species\* and Canopy Position**

**Upper Layer Lifeform**

- Herbaceous
- Shrub
- Tree

**Fuel Model** no data

**Structure Data (for upper layer lifeform)**

	Min	Max
Cover	%	%
Height	no data	no data
Tree Size Class	no data	

Upper layer lifeform differs from dominant lifeform.  
Height and cover of dominant lifeform are:

**Disturbances**

**Non-Fire Disturbances Modeled**

- Insects/Disease
- Wind/Weather/Stress
- Native Grazing
- Competition
- Other:
- Other:

**Fire Regime Group: 5**

I: 0-35 year frequency, low and mixed severity  
II: 0-35 year frequency, replacement severity  
III: 35-200 year frequency, low and mixed severity  
IV: 35-200 year frequency, replacement severity  
V: 200+ year frequency, replacement severity

\*Dominant and Indicator Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

**Historical Fire Size (acres)**

Avg: 100  
Min: 10  
Max:1000

**Fire Intervals (FI):**

Fire interval is expressed in years for each fire severity class and for all types of fire combined (All Fires). Average FI is the central tendency modeled. Minimum and maximum show the relative range of fire intervals, if known. Probability is the inverse of fire interval in years and is used in reference condition modeling. Percent of all fires is the percent of all fires in that severity class. All values are estimates and not precise.

**Sources of Fire Regime Data**

- Literature
- Local Data
- Expert Estimate

	Avg FI	Min FI	Max FI	Probability	Percent of All Fires
<i>Replacement</i>	1000			0.001	41
<i>Mixed</i>					
<i>Surface</i>	714			0.00140	58
<i>All Fires</i>	416			0.00241	

**References**

Brown, James K.; Smith, Jane Kapler. Eds. 2000. Wildland fire in ecosystems: effects of fire on flora. Gen.Tech. Rep. RMRS-GTR-42-vol. 2. Ogden, UT: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. 257 p.

Heineke, Thomas E. 1987. The flora and plant communities of the middle Mississippi river valley. PhD dissertation. S.Illinois University. 653 p.

Kuchler, A.W. 1964. Potential natural vegetation of the conterminous United States. American geographical society. New York, NY. 116 p.

Schmidt, Kirsten M., Menakis, James P., Hardy, Colin C., Hann, Wendell J., Bunnell, David L. 2002. Development of coarse-scale spatial data for wildland fire and fuel management. Gen. Tech. Rep. RMRS-GTR-87. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. 41 p. +CD.

Tingle, John L., Charles V. Klimas, and Thomas L. Foti. 1995. Application of GLO survey notes to bottomland ecosystem management and restoration in the lower Mississippi valley-an example from Desha county Arkansas. IN: Bottomland hardwoods of the Mississippi alluvial valley: characteristics and management of natural function, structure, and composition. Gen. Tech. Rep. GTR-SRS-42. Fayetteville, AR. U.S. Department of Agriculture, Forest Service, Southern Research Station.